Student Name Date Instructor's Name	Student Name Date	Instructor's Name
-------------------------------------	-------------------	-------------------

## PERFORMANCE CHECKLIST 22-9B

## TRACHEOSTOMY CARE

		S	U	Comments
1.	Refer to standard pre-procedure steps 1-9			
2.	Assemble equipment			
3.	Check patient's tracheostomy for sanguineous exudate, edema, and respiratory obstruction			
4.	Perform suctioning if it is needed before performing tracheostomy care			
5.	Position patient in semi-Fowler's position			
6.	Perform hand hygiene			
7.	Position self at head of bed facing patient			
<ul><li>8.</li><li>9.</li></ul>	Don clean glove on nondominant hand; remove old dressing from around tracheostomy stoma, and discard it in appropriate receptacle  Prepare equipment and supplies on overbed table	٥		
	a. Open tracheostomy cleaning kit with aseptic technique	٥		
	b. If basins are packed with sterile gloves, apply one sterile glove to dominant hand; separate basins with dominant hand; use nondominant hand to pour cleansing solution (hydrogen peroxide) in one basin and rinsing solution (normal saline) in another basin	0		
10.	With nondominant hand, unlock and remove inner cannula; place in hydrogen peroxide cleansing solution; never remove outer cannula—if it is expelled by patient, use hemostat to			
	hold tracheostomy open, and call for assistance			

			S	U	Comments
11.		ply second sterile glove, or apply new pair sterile gloves; clean inner cannula			
	a.	Use brush to clean inside and outside of inner cannula			
	b.	Place inner cannula in sterile normal saline solution; rinse thoroughly			
	c.	Inspect inner and outer areas of inner cannula; remove excess liquid			
	d.	Insert inner cannula and lock in place			
12.	out cot usi	ean skin around tracheostomy and tabs of ter cannula with hydrogen peroxide and ton-tipped swabs; clean away from opening ong wipes that are free of lint around the cheostomy opening			
13.	fro pla	may be necessary to rinse cleansing solution m skin; if so, use sterile 4 × 4 gauze pads; ce dry, sterile dressing around tracheostomy eplate		<u> </u>	
14.	Ch	ange cotton tapes			
	a.	If assistance is not available, thread clean tie through opening in flange of outer cannula alongside old tie; if assistance is available, untie one side of cotton tape from outer cannula and replace with clean one			
	b.	Bring clean tape under back of neck			
	c.	If assistance is not available, thread tie through opening in opposite flange of outer cannula alongside old tie; if assistance is available, remove other side from outer cannula and replace with clean tape		0	
	d.	Tie ends of two clean cotton tapes together and position knot appropriately	0		
15.		scultate lung sounds; monitor oxygen uration			

		S	U	Comments
16.	Provide mouth care			
17.	Place paper and pencil within easy reach of patient	<u> </u>		
18.	Document per facility policy			
19.	Refer to standard post-procedure steps 12-17			